

# A DERIVATION OF THREE-DIMENSIONAL INERTIAL TRANSFORMATIONS

**F. Goy<sup>1</sup>**

*Dipartimento di Fisica, Università di Bari  
Via G. Amendola, 173  
I-70126 Bari, Italy  
E-mail: goy@axpba1.ba.infn.it*

June 9, 1997

The derivation of the transformations between inertial frames made by Mansouri and Sexl is generalised to three dimensions for an arbitrary direction of the velocity. Assuming length contraction and time dilation to have their relativistic values, a set of transformations kinematically equivalent to special relativity is obtained. The “clock hypothesis” allows the derivation to be extended to accelerated systems. A theory of inertial transformations maintaining absolute simultaneity is shown to be the only one logically consistent with accelerated movements. Algebraic properties of these transformations are discussed.

Key words: special relativity, synchronization, one-way velocity of light, ether.

## 1 INTRODUCTION

In the last two decades, there has been a revival of ether theories. There are at least two important reasons for this new interest in old ideas:

- The discovery of the microwave background radiation enables us to define a privileged frame in which the temperature of this radiation is isotropic.
- The development of the parametric test theories of special relativity of Mansouri and Sexl [1], and that of Robertson [2], are based

---

<sup>1</sup>Financial support of the Swiss National Science Foundation and the Swiss Academy of Engineering Sciences.

on the assumption that there is at least one inertial frame in which light behaves isotropically. It has shown, among other things, that if the parameters of length contraction and time dilation are well-testable quantities, the parameter corresponding to the synchronization of distant clocks in an inertial frame is not a testable quantity [3]. Consequently, one obtains a set of theories equivalent to special relativity, at least in inertial frames. The use of the so-called “absolute” synchronization, which preserves simultaneity, leads to an ether theory in the sense that it singles out a privileged inertial frame, in which the one-way velocity of light is  $c$  in all directions. In all other frames, the one-way velocity of light is generally direction-dependent; only the two-way velocity of light is constant in all directions and for all frames. This theory of inertial transformations (IT) explains the Michelson-Morley experiment and many others (Römer, stellar aberration [4], Fizeau, Kennedy-Thorndike, Ives-Stillwell, Mössbauer rotor [5], Brillet-Hall [6], Hils-Hall [7], etc...) just as well as SRT.

In their article of 1977, Mansouri and Sexl derived transformations between two inertial frames for an arbitrary synchronization based on two main assumptions: a clock moving at velocity  $v$  relative to the privileged frame slows down its rate according to the relativistic factor  $\sqrt{1 - v^2/c^2}$ , and a moving measuring rod contracts longitudinally with the same factor. Their derivation is done in the case in which the velocity is parallel to the  $X$ -axis, but they give a general formula for an arbitrary direction of the velocity without justifying it. In particular, it is not clear in their article, to what axes the coordinates are referred. The extension to an arbitrary direction of the velocity is not trivial, since in special relativity the product of two boosts with non-collinear velocity is not a boost, but it is the case for parallel velocities. Moreover, because of the Lorentz contraction it is easy to show that an orthogonal system of axes moving with constant velocity relative to an inertial frame appear to be non-orthogonal if the velocity of the moving system is not parallel to one of the axes.

In sections 2, 3 and 4 we generalise the derivation made by Mansouri and Sexl to an arbitrary direction of the velocity. In section 5, we show that only the inertial transformations are consistent with the clock hypothesis when accelerations are involved. In sections 6, analysing basic algebraic properties of the inertial transformations, we discuss the question of the Thomas precession in the context of the inertial theory. In section 7, we discuss the advantages and disadvantages of the inertial theory as well as its domain of application.

## 2 DERIVATION IN THE GENERAL CASE

Following Robertson, and Mansouri and Sexl, we suppose that there is at least one privileged inertial frame  $\Sigma$  in which light behaves isotropically and in which clocks are consequently synchronized with Einstein’s procedure [8, 1]. The length of an object having an arbitrary motion in  $\Sigma$  is

defined, as usual, as the distance measured with unit rods at rest in  $\Sigma$  between the simultaneous (in  $\Sigma$ ) position of the endpoints of the object. Without loss of generality, we can choose an orthonormal set  $\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3$  in  $\Sigma$  to which the coordinates of a point are referred. By “vector” we shall mean, in the following, not a triplet of numbers, but a geometrical object, representing a physical one and independent of the system of coordinates. We suppose further that space is homogeneous and isotropic and that time is homogeneous. Another inertial system  $S$  is given system of axes whose origin coincides with the origin of  $\Sigma$  at time  $T = 0$  of  $\Sigma$  and which we shall not specify further for the time being. If we write the coordinates of a point in  $\Sigma$  as  $(X^\alpha, T)$  and as  $(x^\alpha, t)$  in  $S$  ( $\alpha = 1, 2, 3$ ), then the conditions on the origin, and of homogeneity and isotropy, imply that the relations between the coordinates are linear [9]. We can write:

$$\begin{aligned} x^\alpha &= a^\alpha_\beta X^\beta + b^\alpha T \\ t &= s_\alpha X^\alpha + a^0_0 T \end{aligned} \quad (1)$$

where repeated indices are summed over 1, 2, 3 and the coefficients  $a^\alpha_\beta$ ,  $b^\alpha$ ,  $s_\alpha$  and  $a^0_0$  do not depend on the coordinates. The matrix  $a^\alpha_\beta$  contains information about longitudinal and transversal length contraction factors, as well as on the choice of the system of axes in  $S$ .  $a^0_0$  is related to time dilation. The parameters  $b^\alpha$  describe the velocity of the origin of the axes of  $S$  and the  $s_\alpha$  are parameters characterising the synchronization of clocks in  $S$ . Indices are formally lowered and raised with  $\text{diag}(1,1,1)$ : for example:  $s^\alpha = \delta^{\alpha\beta} s_\beta$ , so that covariant and contravariant components have numerically the same value.

We suppose further that the origin of  $S$  is moving with a velocity  $\mathbf{v} = v^\alpha \mathbf{E}_\alpha$ . Imposing this condition in (1) we obtain:

$$\begin{aligned} x^\alpha &= a^\alpha_\beta (X^\beta - v^\beta T) \\ t &= s_\alpha X^\alpha + a^0_0 T \end{aligned} \quad (2)$$

We suppose now that *bodies moving through  $\Sigma$  are contracted parallel to their motion in  $\Sigma$  by the relativistic factor  $R = \sqrt{1 - v^2/c^2}$* . In order to express it mathematically, we choose a set  $(\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3)$  of vectors coincident with  $(\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3)$  at  $T = 0$  if  $\mathbf{v} = \mathbf{0}$ . Because accelerations *per se* have no effects of contraction on ideal rods [10, p. 256], the system chosen is in fact the system we would obtain after acceleration up to velocity  $\mathbf{v}$  if the two systems coincided originally. We can decompose the vectors  $\mathbf{E}_\alpha$  in a part  $\mathbf{E}_{\alpha\parallel}$  parallel to  $\mathbf{v}$  and a part  $\mathbf{E}_{\alpha\perp}$  perpendicular to  $\mathbf{v}$ :

$$\mathbf{E}_{\alpha\parallel} = (\hat{\mathbf{v}} \cdot \mathbf{E}_\alpha) \hat{\mathbf{v}} \quad (3)$$

and

$$\mathbf{E}_{\alpha\perp} = \mathbf{E}_\alpha - (\hat{\mathbf{v}} \cdot \mathbf{E}_\alpha) \hat{\mathbf{v}} \quad (4)$$

where  $\hat{\mathbf{v}} = \mathbf{v}/|\mathbf{v}|$ . So our last assumption is equivalent to:

$$\begin{aligned}\mathbf{e}_\alpha &= R\mathbf{E}_{\alpha\parallel} + \mathbf{E}_{\alpha\perp} \\ &= [\delta_\alpha^\beta + (R-1)\hat{v}_\alpha\hat{v}^\beta]\mathbf{E}_\beta\end{aligned}\quad (5)$$




Figure 1: Configuration of the axes in two dimensions for  $R = 1/2$  and the velocity  $\mathbf{v}$  making an angle of  $30^\circ$  with  $\mathbf{E}_1$ .

This last equation gives us the relation between the vectors of the two bases. From (5) and (2), we obtain the transformations of the coordinates referred to these axes, by expressing the position of a point in the two bases:

$$\begin{aligned}x^\alpha &= [\delta_\beta^\alpha + (\frac{1}{R} - 1)\hat{v}^\alpha\hat{v}_\beta]X^\beta - \frac{1}{R}v^\alpha T \\ t &= s_\alpha X^\alpha + a_0^0 T\end{aligned}\quad (6)$$

The last equation of (6) can be rewritten as:

$$t = s_\alpha(X^\alpha - v^\alpha T) + (s_\alpha v^\alpha + a_0^0)T \quad (7)$$

We now suppose that *a clock identical to the clocks in  $\Sigma$  is at the origin of  $S$  and is slowed down by a factor  $R$* . We obtain from (7) for the time coordinate difference that:

$$\Delta t = (s_\alpha v^\alpha + a_0^0)\Delta T = R\Delta T \quad , \quad (8)$$

so that (6) can now be written as:

$$\begin{aligned}x^\alpha &= [\delta_\beta^\alpha + (\frac{1}{R} - 1)\hat{v}^\alpha\hat{v}_\beta]X^\beta - \frac{1}{R}v^\alpha T \\ t &= s_\alpha(X^\alpha - v^\alpha T) + RT\end{aligned}\quad (9)$$

Only the tree parameters  $s_\alpha$  characterising the synchronization in  $S$  remain unknown. The inverse of (9) is given by:

$$\begin{aligned} X^\alpha &= [\delta_\beta^\alpha + (R - \mathbf{s} \cdot \mathbf{v})(1 - \frac{1}{R})\hat{v}^\alpha \hat{v}_\beta - v^\alpha s_\beta]x^\beta + \frac{v^\alpha}{R}t \\ T &= [(\frac{1}{R} - 1)\hat{v}_\alpha(\mathbf{s} \cdot \hat{\mathbf{v}}) - \frac{s_\alpha}{R}]x^\alpha + \frac{1}{R}t, \end{aligned} \quad (10)$$

where  $\mathbf{s} = s^\alpha \mathbf{E}_\alpha$ . In (9) and (10), the inertial transformations are given by  $\mathbf{s} = \mathbf{0}$ , while the Lorentz transformations are given by  $\mathbf{s} = -\mathbf{v}/c^2 R$ .

### 3 ONE-WAY VELOCITY OF LIGHT IN S

Given the isotropy of the velocity of light in  $\Sigma$ , two events which are light-like obey the equation:

$$\Delta X^\alpha \Delta X_\alpha - c^2 \Delta T^2 = 0, \quad (11)$$

where  $\Delta$  expresses the difference between the coordinates of the two events. In order to calculate the left-hand-side of (11) in  $S$ , we define new parameters  $h_\alpha$  characterising the synchronization in  $S$ , which are more convenient than the  $s_\alpha$  for this calculation [11]:

$$\mathbf{h} = -(cR\mathbf{s} + \frac{\mathbf{v}}{c}) \quad (12)$$

The Lorentz transformations are obtained for  $\mathbf{h} = \mathbf{0}$ , while the inertial transformations are obtained for  $\mathbf{h} = -\mathbf{v}/c$ . If we have synchronized the clocks with  $\mathbf{h} = \mathbf{0}$  in  $S$ , we can obtain an “absolute” synchronization  $\mathbf{h} = -\mathbf{v}/c$  by performing a resynchronization of our clocks. For a general  $\mathbf{h}$  the resynchronization is given by:

$$ct^{\mathbf{h}} = ct^{\mathbf{h}=\mathbf{0}} - \mathbf{h} \cdot \mathbf{x} \quad (13)$$

For the calculation of (11), we can first use a Lorentz boost to reach  $S$ . We obtain:

$$\Delta x^\alpha \Delta x_\alpha - c^2 \Delta(t^{\mathbf{h}=\mathbf{0}})^2 = 0, \quad (14)$$

because the Lorentz transformations leaves the Minkowski metric invariant. Resynchronizing the clocks in  $S$  with (13), we obtain the equation :

$$c^2 \Delta t^2 + 2c \Delta t \mathbf{h} \cdot \Delta \mathbf{x} + (\mathbf{h} \cdot \Delta \mathbf{x})^2 - (\Delta \mathbf{x})^2 = 0, \quad (15)$$

which can be written with the metric:

$$g_{\alpha\beta} = \begin{pmatrix} -1 & -h_1 & -h_2 & -h_3 \\ -h_1 & 1 - h_1^2 & -h_1 h_2 & -h_1 h_3 \\ -h_2 & -h_1 h_2 & 1 - h_2^2 & -h_2 h_3 \\ -h_3 & -h_1 h_3 & -h_2 h_3 & 1 - h_3^2 \end{pmatrix} \quad (16)$$

From (15), the velocity of light in  $S$  can easily be calculated and is given in the direction  $\hat{\mathbf{n}} = \Delta x^\alpha \mathbf{e}_\alpha / \|\mathbf{x}\|$  by:

$$\mathbf{c}(\hat{\mathbf{n}}) = \frac{c\hat{\mathbf{n}}}{1 - \mathbf{h} \cdot \hat{\mathbf{n}}} \quad (17)$$

First, note that the one-way velocity of light is generally direction dependent in  $S$ . The two-way velocity of light is on the other hand constant in all directions. The inverse of the two-way velocity of light  $\overleftrightarrow{\mathbf{c}}(\hat{\mathbf{n}})$  in direction  $\hat{\mathbf{n}}$  is given by:

$$\frac{1}{\|\overleftrightarrow{\mathbf{c}}(\hat{\mathbf{n}})\|} = \frac{1}{2} \left( \frac{1}{\|\mathbf{c}(\hat{\mathbf{n}})\|} + \frac{1}{\|\mathbf{c}(-\hat{\mathbf{n}})\|} \right) = \frac{1}{c} \quad , \quad (18)$$

in all directions. So we see that assuming the length contraction and time dilation parameters to have their relativistic value implies a two-way velocity of light constant in all directions. Other derivations had already shown that, assuming the time dilation parameter to have its relativistic value, and assuming also the constancy of the two-way velocity of light, the length contraction parameters have their relativistic values [13].

Secondly, notice that the norm of a vector  $\mathbf{x}$  in  $S$ , whose coordinates are  $x^\alpha$  ( $\alpha = 1, 2, 3$ ) is given by  $\|\mathbf{x}\| = \sqrt{x^\alpha x_\alpha}$  despite the fact that, seen from  $\Sigma$ , the set  $\mathbf{e}_\alpha$  ( $\alpha = 1, 2, 3$ ) is not orthonormal. But in  $S$ , lengths are measured with unit rods at rest in  $S$ . These measuring instruments as well as the objects they measure are contracted relative to  $\Sigma$ , so that the set  $\mathbf{e}_\alpha$  ( $\alpha = 1, 2, 3$ ) is found to be orthonormal in  $S$  and consequently the norm of a vector is calculated with a diagonal 3-dimensional metric. A mathematical proof of this last remark can be given making use of (16). As is well known, in the formalism of general relativity, which includes transformations (9) as a special case, the spatial part  $\gamma_{ij}$  of the metric of a four dimensional space-time metric is not only given by the space-space coefficients of this metric but by the expression:

$$\gamma_{ij} = \left( g_{ij} - \frac{g_{i0}g_{j0}}{g_{00}} \right) \quad i, j = 1, 2, 3 \quad , \quad (19)$$

which gives, in the case of (16),  $\text{diag}(1,1,1)$ ; that is, an orthonormal metric in Cartesian coordinates. Eq. (19) corresponds to the fact that distances are measured with rigid measuring rods (see [10, p. 269]), which can be defined in terms of two-way light signals [11].

Note also that in (16), the components of the vector potential (or gravitomagnetic potential) ( $g_{01}, g_{02}, g_{03}$ ) are non zero. The gravitomagnetic field [12] is defined as  $\mathbf{H} = \nabla \wedge (-\mathbf{h})$ . As in the case of the Minkowski metric, we have from (16):  $\mathbf{H} = \mathbf{0}$ , whenever  $(-\mathbf{h})$  can be written in the form  $-\mathbf{h} = \nabla f$ , where  $f$  is a function.

## 4 SYNCHRONISATION WITH ONE PARAMETER

If we now assume that *there is in  $S$  only one privileged direction given by the velocity  $\mathbf{v}$* , we can restrict the three parameters  $s_\alpha$  to one parameter. We suppose that the one-way velocity of light has the same norm for any direction making the same angle with  $\mathbf{v}$ . This means that  $\hat{\mathbf{n}}_1 \cdot \mathbf{h} = \hat{\mathbf{n}}_2 \cdot \mathbf{h} \forall \hat{\mathbf{n}}_1, \hat{\mathbf{n}}_2$ , such that  $\hat{\mathbf{n}}_1 \cdot \mathbf{v} = \hat{\mathbf{n}}_2 \cdot \mathbf{v}$ . For  $\mathbf{h} = \mathbf{0}$ , this condition is trivially fulfilled, otherwise it is equivalent to  $\mathbf{h} = h\hat{\mathbf{v}}$ , so that also  $\mathbf{s} = s\hat{\mathbf{v}}$ . We remain with only one parameter characterising the synchronization and equation (9) becomes:

$$\begin{aligned} x^\alpha &= [\delta_\beta^\alpha + (\frac{1}{R} - 1)\hat{v}^\alpha \hat{v}_\beta] X^\beta - \frac{1}{R} v^\alpha T \\ t &= s \hat{v}_\alpha X^\alpha + (R - vs) T \end{aligned} \quad (20)$$

with its inverse given by:

$$\begin{aligned} X^\alpha &= [\delta_\beta^\alpha + (R - vs - 1)\hat{v}^\alpha \hat{v}_\beta] x^\beta + \frac{v^\alpha}{R} t \\ T &= -s(\hat{v}_\alpha x^\alpha) + \frac{1}{R} t \end{aligned} \quad (21)$$

Substituing  $s = -v/c^2 R$  in (20), we obtain a Lorentz-boost in (3+1) dimension (see [15, eqs:(2.1.20),(2.1.21)]):

$$\begin{aligned} x^\alpha &= [\delta_\beta^\alpha + (\frac{1}{R} - 1)\hat{v}^\alpha \hat{v}_\beta] X^\beta - \frac{1}{R} v^\alpha T \\ t &= \frac{1}{R} \left( T - \frac{v_\alpha X^\alpha}{c^2} \right) \end{aligned} \quad (22)$$

Substituing  $s = 0$ , we obtain the (3+1) dimensional inertial transformation:

$$\begin{aligned} x^\alpha &= [\delta_\beta^\alpha + (\frac{1}{R} - 1)\hat{v}^\alpha \hat{v}_\beta] X^\beta - \frac{1}{R} v^\alpha T \\ t &= R T \end{aligned} \quad (23)$$

In the following table, we show also for which value of  $h$  the Lorentz transformation and the inertial transformation are recovered.

|     | LT                 | IT             |
|-----|--------------------|----------------|
| $h$ | 0                  | $-\frac{v}{c}$ |
| $s$ | $-\frac{v}{c^2 R}$ | 0              |

Different choices of  $s$  represent different conventions of synchronization of distant clocks, corresponding to different values of the one-way speed of light. In theories in which no interaction between the measuring apparatus

and the system is assumed, the setting of clocks cannot change the objective nature of physical processes, but only our description of it. Thus, the transformations (22) and (23) represent the same transformation of spacetime, but relative to different spacetime coordinates resulting from the choice of different methods of synchronization of clocks. The above remark is not restricted to the two particular values of  $s$  in equations (22) and (23) and we can say that (20) represents a set of transformations equivalent to the transformation of Lorentz between  $\Sigma$  and  $S$ . Strictly speaking, this assertion is true in inertial systems only, since our derivation of (20) was made in this class of systems. Different coordinatisations than that of the Lorentz transformation are obviously equivalent: nobody claims, for example, that we obtain different physical results if we use three axes which are not orthonormal. The intrasystemic synchronization of distant clocks, by defining the simultaneity of distant events, allows the attribution of a common time to all points of an inertial system and thus corresponds to the choice of an axis in the spacetime domain. Without synchronizing clocks time would not even be defined, but we would only have a “time” for each clock with no connection between them and thus no way of attributing a time coordinate to an event. In relativity and related theories, clocks are synchronized by means of light signals whose one-way velocity is *assumed* as stressed by Einstein in 1916 [16]: *“that light requires the same time to traverse the path AM...as the path BM [M being the midpoint of the line AB] is in reality neither a supposition nor a hypothesis about the physical nature of light, but a stipulation which I can make of my own free will”*. Numerous authors have stressed the conventional nature of the one-way speed of light (for a review see: [17]). The explanation of Anderson and Stedman is of particular interest [11]: *“In order to synchronize spatially separated clocks, signals must be sent between them and the time of passage of such signals must be known. A maximum specification will only arise through the use of light signals by virtue of their property of being first signals. But we are caught in a circle here as the time of passage for first signals can only be obtained by prior knowledge of the synchronization of clocks. The essence of the “conventionalist” position is that this circularity is inescapable and that no fact of nature permits a unique determination of either the simultaneity relation within an inertial frame or the speed of light in a given direction”*. So the freedom of choice of the synchronization of clocks is equivalent to the freedom of choice in the assumption of the value of the one-way speed of light, which leads to a freedom in the choice of the spacetime coordinate system. From this point of view, it is obvious that (20) represents a set of transformations experimentally equivalent to the Lorentz transformations between  $\Sigma$  and  $S$ .

## 5 THE CLOCK HYPOTHESIS

In this section, we will show that in the case of accelerated systems, only the transformation with  $s = 0$  is consistent with the “clock hypothesis”.



In this sense of logical consistency, transformations (22) and (23) are no more equivalent when extended to accelerated systems.

The “clock hypothesis” states that the rate of an ideal clock accelerated relative to an inertial frame is identical to the rate of a similar clock in the instantaneously comoving inertial frame. With other words, the rate of clocks is not influenced by accelerations *per se*, when seen from inertial frames. It also supposed that real clocks exist in nature, which approach the conditions of the clock hypothesis. To our knowledge, this assumption was first implicitly used by Einstein in 1905 [8] and was superbly confirmed in the CERN muon storage ring experiment [14], where the muons had a time of decay depending only on their velocity (in agreement with the time dilation formula) despite the fact that their acceleration was of  $10^{18}g$ . We stress here that the clock hypothesis is an assumption logically independent of the assumptions used to derive (20), and of the two postulates of special relativity.

Using the clock hypothesis in order to extend the theory to the possibility of temporary acceleration, we will restrict the last unknown parameter. The necessity of this extension is obvious, since a frame needs to accelerate to go from an uniform velocity to an other uniform velocity. Developing an argument of Bell [18] and Selleri [13], we imagine the following situation. Two clocks are in two spaceships at rest in  $\Sigma$ . The first clock A is at the origin of  $\Sigma$  and the second clock B at a point P, whose position is given by a vector  $\mathbf{d}$ . At  $T = 0$ , the two spaceships begin to accelerate in straight line under the constraint of a constant force equal for both spaceships. They continue their acceleration up to a time  $\bar{T}$ , and then continue in uniform motion. In  $\Sigma$  the law of motion of relativity is valid because clocks are synchronized with an Einstein’s procedure. It follows that the two spaceships will have an hyperbolic motion whose equation is given for the time  $T \leq \bar{T}$  by:

$$\begin{aligned}\mathbf{X}_A(T) &= \frac{\hat{\mathbf{n}}c^2}{a} \left\{ \left[ 1 + \left( \frac{aT}{c} \right)^2 \right]^{\frac{1}{2}} - 1 \right\} \\ \mathbf{X}_B(T) &= \frac{\hat{\mathbf{n}}c^2}{a} \left\{ \left[ 1 + \left( \frac{aT}{c} \right)^2 \right]^{\frac{1}{2}} - 1 \right\} + \mathbf{d} \quad ,\end{aligned}\tag{24}$$

where  $\hat{\mathbf{n}}$  is a unit vector characterising the direction of propagation and  $a$  is the ratio of the force by the rest mass of the spaceship. For  $T > \bar{T}$ , the same term  $\hat{\mathbf{n}}v(T - \bar{T})$  ( $v$  is the velocity attained at  $T = \bar{T}$ ) must be added to both equations (24) so that  $\forall T : \mathbf{X}_B(T) - \mathbf{X}_A(T) = \mathbf{X}_B(0) - \mathbf{X}_A(0) = \mathbf{d}$ . That is the distance between the spaceships remains constant in  $\Sigma$ . The velocity of the spaceships for  $0 \leq T \leq \bar{T}$  is given by:

$$\dot{\mathbf{X}}_{A,B}(T) = \frac{\hat{\mathbf{n}}aT}{\left[ 1 + \left( \frac{aT}{c} \right)^2 \right]^{\frac{1}{2}}}\tag{25}$$

The clock hypothesis states mathematically that:

$$dt = \left( 1 - \frac{\dot{X}_{A,B}^2(T)}{c^2} \right)^{\frac{1}{2}} dT \quad , \quad (26)$$

where  $t$  is the time coordinate in the accelerated system  $S$ . Integrating (26) with (25) and the given initial conditions, we obtain for the time of  $S$  as function of the time of  $\Sigma$ , for  $0 \leq T \leq \bar{T}$ :

$$t_A(T) = t_B(T) = \frac{c}{a} \operatorname{arcsinh}\left(\frac{aT}{c}\right) \quad (27)$$

and for  $T > \bar{T}$ :

$$t_A(T) = t_B(T) = \frac{c}{a} \operatorname{arcsinh}\left(\frac{a\bar{T}}{c}\right) + \sqrt{1 - v^2/c^2} (T - \bar{T}) \quad (28)$$

so that  $\forall T : t_A(T) - t_B(T) = 0$ . Thus we see that though slowing their rate, clocks A and B maintain the simultaneity of events.

On the other hand, we can calculate the difference of time coordinates in  $S$  using directly an inertial transformation with a general  $s$ . Using (20), we find for two simultaneous events in  $\Sigma$

$$t_B^s(T) - t_A^s(T) = s \hat{\mathbf{v}} \cdot (\mathbf{X}_B(T) - \mathbf{X}_A(T)) = s \hat{\mathbf{v}} \cdot \mathbf{d} \quad (29)$$

Since this last equation must hold for all  $\mathbf{d}$ , we obtain  $s = 0$  comparing with (27). In particular, an observer moving with the spaceships would have to resynchronize the clocks in order to obtain the Lorentz-transformations between his frame and  $\Sigma$  [19]. Nevertheless, such a resynchronization allows to define a time coordinate only locally as explained by Misner, Thorne and Wheeler [20]: *“Difficulties also occur when one considers an observer who begins at rest in one frame, is accelerated for a time, and maintains thereafter a constant velocity, at rest in some other inertial coordinate system. Do his motion define in any natural way a coordinate system? Then this coordinate system (1) should be the inertial frame  $x^\mu$  in which he was at rest for time  $x^0$  less than 0, and (2) should be the other inertial frame  $x^{\mu'}$  for times  $x^{0'} > T'$  in which he was at rest in the other frame. Evidently some further thinking would be required to decide how to define the coordinates in the regions not determined by these two conditions. More serious, however, is the fact that these two conditions are inconsistent for a region of spacetime that satisfies simultaneously  $x^0 < 0$  and  $x^{0'} > T'$ .”* Further they conclude that the problem possess a solution only in the immediate vicinity of the observer. With other words special relativity is valid only locally when accelerations come into play. That is indeed one of the reasons why special relativity is valid only locally within the frame of general relativity. The reader thinks perhaps that the problem

above does not occur in general relativity, where there is absolutely no doubt that accelerated observers can be considered. General relativity is coordinate independent and thus any synchronization can be used. The latter statement is true only locally, because some synchronizations allow a global definition of time, others (in particular Einstein's synchronization) are path dependent when the metric is non-static [21][10, p. 379]. This implies in particular that the binary relation between two events "A is synchronous with B" is not an equivalence relation in the latter case. Applying the formalism of general relativity to the rotating disk, one can show that only a synchronization corresponding to  $s = 0$  allows a global and self-consistent definition of time on the disk [22]. Synchronization problems in the frame of general relativity is beyond the scope of this paper. We have mentioned them shortly in order to show that the problem expounded in this section does not disappear in general relativity.

Coming back to the problem of two accelerated clocks treated in this section we conclude that the only theory whose synchronization of clocks is *logically consistent* with the clock hypothesis is the inertial theory using  $s = 0$ . With other word, the natural behaviour of clocks (that is without resynchronisation by an observer) is described by a theory using an "absolute" simultaneity ( $s = 0$ ). Moreover a time defined in this way is global as it should be.

This result could have been derived with weaker assumptions than the clock hypothesis. For example, we would have obtained the same result if the rate of the clocks were dependent on their accelerations (seen from an inertial frame). What is important in the above reasoning is that clocks A and B, which are Einstein's synchronised in  $\Sigma$ , are submitted to the same influences, so they have the same rate, so maintain the simultaneity of  $\Sigma$ . This idea was already expressed by Einstein in his article of 1907 where he first used the principle of equivalence and discussed the behaviour of two clocks in an accelerated frame  $\Sigma$  which was at rest in an inertial frame  $S$  at time  $t = 0$  (the clocks are Einstein's synchronised in  $S$ ). He wrote [23] *"Indeed, being given that two arbitrary clocks of  $\Sigma$  are synchronous at time  $t = 0$  with respect to  $S$  and that they are subject to the same motions, they remain constantly synchronous with respect to  $S$ . But for this reason, and according to the paragraph 4, [that is according to the Lorentz transformations] they do not work in a synchronous way with respect to a system  $S'$  momentarily at rest with respect to  $\Sigma$  but in motion with respect to  $S$ ."*

## 6 SOME ALGEBRAIC PROPERTIES OF INERTIAL TRANSFORMATIONS

We call  $A(\mathbf{v})$  the inertial transformation (20) between  $\Sigma$  and  $S$ : The transformations  $\Omega(\mathbf{v}, \mathbf{w})$  between two inertial systems  $S$  and  $S'$  having a velocity  $\mathbf{v}$  and  $\mathbf{w}$  relative to  $\Sigma$  can be found by elimination of the coordinates of

the  $\Sigma$  system as:

$$\Omega(\mathbf{v}, \mathbf{w}) = A(\mathbf{v})A^{-1}(\mathbf{w}) \quad (30)$$

and have in general a complicated form. Some of the algebraic properties of the set of all transformations  $\Omega(\mathbf{v}, \mathbf{w})$  can be demonstrated using the definition (30). Notice that each element  $\Omega(\mathbf{v}, \mathbf{w})$  is parametrized by 6 parameters  $v^\alpha, w^\alpha$ , which can vary continuously, subject only to the conditions  $\|\mathbf{v}\| \leq c$  and  $\|\mathbf{w}\| \leq c$ .

It does not make physically sense to multiply  $\Omega(\mathbf{v}, \mathbf{w}) : S \rightarrow S'$  with  $\Omega(\mathbf{u}, \mathbf{r}) : S'' \rightarrow S'''$ . Mathematically, we can multiply two such transformations. Using (30), we obtain:

$$\Omega(\mathbf{v}, \mathbf{w})\Omega(\mathbf{u}, \mathbf{r}) = A(\mathbf{v})A^{-1}(\mathbf{w})A(\mathbf{u})A^{-1}(\mathbf{r}) \quad , \quad (31)$$

which is in general not a transformation of type  $\Omega$ , but an object depending on 12 independant parameters. So the set of all transformations  $\Omega$ , with the composition law defined above, *does not form a group* because it is not closed. What makes physically sense is to go from  $S \rightarrow S'$  and then from  $S' \rightarrow S''$ . In this case, the composition of two transformations  $\Omega(\mathbf{v}, \mathbf{w}) : S \rightarrow S'$  and  $\Omega(\mathbf{w}, \mathbf{u}) : S' \rightarrow S''$ :

$$\Omega(\mathbf{v}, \mathbf{w})\Omega(\mathbf{w}, \mathbf{u}) = A(\mathbf{v})A^{-1}(\mathbf{w})A(\mathbf{w})A^{-1}(\mathbf{u}) = \Omega(\mathbf{v}, \mathbf{u}) \quad , \quad (32)$$

gives again a transformation of type  $\Omega$ :  $\Omega(\mathbf{v}, \mathbf{u})$  from  $S$  to  $S''$ . The properties of the inertial transformations are similar to those of evolution operators. Despite of the fact that the  $\Omega$ 's do not form a group, for the cases which are physically relevant, everything is well defined. We have neutral elements, every element has an inverse and the composition is associative:

1.  $\Omega(\mathbf{v}, \mathbf{v}) = I \quad \forall \mathbf{v}$
2.  $\Omega(\mathbf{v}, \mathbf{w})\Omega(\mathbf{w}, \mathbf{v}) = I \quad \forall \mathbf{v}, \mathbf{w}$
3.  $[\Omega(\mathbf{v}, \mathbf{w})\Omega(\mathbf{w}, \mathbf{u})]\Omega(\mathbf{u}, \mathbf{r}) = \Omega(\mathbf{v}, \mathbf{w})[\Omega(\mathbf{w}, \mathbf{u})\Omega(\mathbf{u}, \mathbf{r})]$   
 $= \Omega(\mathbf{v}, \mathbf{r}) \quad \forall \mathbf{v}, \mathbf{w}, \mathbf{u}, \mathbf{r}$

Moreover we have  $\det \Omega(\mathbf{v}, \mathbf{w}) = 1$ , whereas the determinant of the spatial part only is given by  $R(v)/R(w)$ . Note that a transformation  $A(\mathbf{v}) : \Sigma \rightarrow S$  can also be expressed with the following formula:

$$R_{\hat{\mathbf{n}}}(\Psi)A(R_{\hat{\mathbf{n}}}(-\Psi)\mathbf{v})R_{\hat{\mathbf{n}}}(-\Psi) = A(\mathbf{v}), \quad (33)$$

where  $\hat{\mathbf{n}}$  is a unit vector and  $R_{\hat{\mathbf{n}}}(\Psi)$  is a rotation around this vector of an angle  $0 \leq \Psi < \pi$ , measured in  $\Sigma$ . As a special case of equation (33), we can express  $A(\mathbf{v})$  as given by (20) in terms of a inertial boost along the  $X$ -axis and a rotation.

So far we have always considered axes of  $\Sigma$  and  $S$  in the positions described in section 2: they would coincide if their relative velocity vanishes. If we want to describe transformations of coordinates in which the axes

$\mathbf{e}_\alpha$  are rotated to new axes  $\mathbf{e}'_\alpha$  such that  $\mathbf{e}'_\alpha = R_{\hat{\mathbf{n}}}(\Psi)\mathbf{e}_\alpha$ , then this more general transformation  $K$  from  $\Sigma$  to  $S$  is the combination of a “boost” and a rotation. We have:

$$K(\mathbf{v}, \hat{\mathbf{n}}, \psi) = R_{\hat{\mathbf{n}}}(-\Psi)A(\mathbf{v}) = A(R_{\hat{\mathbf{n}}}(-\Psi)\mathbf{v})R_{\hat{\mathbf{n}}}(-\Psi) \quad (34)$$

From (34) the corresponding type of transformations between  $S$  and  $S'$ ,  $J$  for axes with arbitrary rotations, will be of the type:

$$J = KK'^{-1} = A(\mathbf{v})R_{\hat{\mathbf{m}}}(\Theta)A^{-1}(\mathbf{w}) \quad (35)$$

We have seen in (32) that the composition of two transformations without rotation,  $\Omega(\mathbf{v}, \mathbf{w}): S \rightarrow S'$  and  $\Omega(\mathbf{w}, \mathbf{u}): S' \rightarrow S''$  gives a transformation  $\Omega(\mathbf{v}, \mathbf{u}): S \rightarrow S''$ , without rotation. This law is valid for any velocities  $\mathbf{v}, \mathbf{w}, \mathbf{u}$ , that is also when the velocity from  $S'$  relative to  $S$  and the velocity of  $S''$  relative to  $S'$  are not collinear. This is a crucial difference with special relativity, where the Lorentz boosts do not form a subgroup of the orthochrone homogeneous proper Lorentz group [24]. In particular, the composition of two Lorentz boosts with non-collinear velocities is not a Lorentz boost, but a Lorentz boost multiplied by a rotation (A). This give rise to Thomas precession (B)[25]. Does it mean that the Thomas precession can not be deduced from the theory of inertial transformations? No because in logic:  $(A \Rightarrow B)$  does not imply  $(\neg A \Rightarrow \neg B)$ . From the absence of rotation in the multiplication of two “inertial boosts”  $(\neg A)$ , one cannot conclude to the absence of Thomas precession  $(\neg B)$ . Other authors than Thomas ([27, 28, 29], see also: [26, ch. 11] and [30]) have given other derivations of the Thomas precession. In their case dynamical effects are explained with dynamical causes and the algebraic properties of the Lorentz transformations do not come directly into play. It could be the subject of a future article to derive the Thomas precession in the frame of the inertial theory. Note that the Thomas precession is in fact well tested in the case of the precession of the muons in the muon storage ring ([31, 27]).

## 7 DISCUSSION

In 1977, Mansouri and Sexl wrote : “*A convention about clock synchronization can be choosen that does maintain absolute simultaneity. Based on this convention an ether theory can be constructed that is, as far as kinematic is concerned (dynamics will be studied in a later paper in this series), equivalent to special relativity*”. Note that this paper was never written, but the Maxwell’s equations were studied by Chang [32] and Rembieliński [33], the dynamics by the latter and Selleri [34]. One can say that from a dynamical point of view this theory is also equivalent to special relativity. The reader is surely asking himself what is the advantage of introducing an ether (privileged frame  $\Sigma$ ) in a theory equivalent to special relativity. Of course this can be done theoretically, but since this theory is equivalent

to special relativity it ensures us also that this ether or the velocity of an inertial frame  $S$  respective to  $\Sigma$  is not detectable within the frame of this theory. Is the development of such theories not a step back in the history of science? I would say that it is a step back when we consider inertial frames only, which allow us to make two steps forward when we consider accelerated frames. We have seen in section 5, that two clocks which are synchronized with an Einstein's procedure in  $\Sigma$  maintain an absolute simultaneity respective to  $\Sigma$  after and during acceleration and thus the theory of inertial transformation is advantaged in the sense that it corresponds to the behaviour of real clocks. This advantage is of logical nature in the sense that the theory of inertial transformation allows a global and self-consistent definition of time, where special relativity cannot do it. This advantage is still present in general relativity where the Einstein's synchronization does not allow to define time globally and self-consistently when the metric is non-static. The difference between the two theories is not really of experimental nature in the sense that we cannot measure the absolute velocity of an inertial frame  $S$  with respect to  $\Sigma$ , when  $\Sigma$  is interpreted as an objectively privileged frame. As already stated by Vetharaniam and Stedman [3], the synchronization of clocks is also conventional in  $\Sigma$ , so that we can call  $\Sigma$  any inertial frame: for any practical purposes  $\Sigma$  is the inertial frame where the clocks are first synchronized with an Einstein's procedure.

## 8 CONCLUSION

A derivation of the transformations between inertial systems was made in the general three-dimensional case. Under the main assumptions that there is at least one inertial frame where light behaves isotropically and that time dilation and length contraction have their relativistic values, we find the whole set of theories kinematically equivalent to special relativity. In the derivation, the notion of vector as an object independent of the coordinates system is used. This approach allows one to see clearly the sort of system of axes to which the coordinates are referred. Moreover, accelerated movements are not *a priori* excluded in this reasoning. The constancy of the two-way velocity of light is shown to be a consequence of the given assumptions. For a general parameter of synchronization, the one-way velocity of light is generally direction dependant and is only equal to  $c$  in the case of special relativity theory. An extension of the transformations to accelerated systems, through the clock hypothesis, shows that only a theory maintaining absolute simultaneity is logically self-consistent.

## 9 ACKNOWLEDGEMENT

I want to thanks the Physics Departement of Bari University for hospitality, and Prof. F. Selleri for its kind suggestions and criticisms.

## REFERENCES

- [1] R. Mansouri and R.U. Sexl, *Gen. Rel. Grav.* **8**, 497–513 (1977); **8**, 515–524 (1977); **8**, 809–813 (1977).
- [2] H.P. Robertson, *Rev. Mod. Phys.* **21**, 378–382 (1949).
- [3] I. Vetharaniam and G.E. Stedman, *Phys. Lett. A* **183**, 349–354 (1993).
- [4] F. Goy, *Found. Phys. Lett.* **9**, 165–174 (1996).
- [5] D.C. Champeney, G.R. Isaak and A.M. Khan, *Phys. Lett.* **7**, 241–243 (1963).
- [6] A. Brillat and J.L. Hall, *Phys. Rev. Lett.* **42**, 549–552 (1979).
- [7] D. Hills and J.L. Hall, *Phys. Rev. Lett.* **64**, 1697–1700 (1990).
- [8] A. Einstein, *Ann. Phys.* **17**, 891–921 (1905).
- [9] F. Selleri, in *Frontiers of Fundamental Physics*, Ed. by M. Barone and F. Selleri (Plenum Press, New-York, 1994), pp. 181–192.
- [10] C. Møller, *The Theory of Relativity*, 2nd ed. (Oxford University Press, Oxford, 1972), p. 379.
- [11] R. Anderson and G.E. Stedman, *Found. Phys. Lett.* **5**, 199–220 (1993).
- [12] I. Ciufolini and J.A. Wheeler, *Gravitation and Inertia* (Princeton University Press, Princeton, 1995).
- [13] F. Selleri, *Phys. Essays* **8**, 342–349 (1995).
- [14] J. Bailey et al., *Nature* **268**, 301–304 (1977).
- [15] S. Weinberg, *Gravitation and Cosmology* (John-Wiley and Sons, New-York, 1972).
- [16] A. Einstein, *Relativity, The Special, The General Theory* (Chicago, 1951) p. 18.
- [17] F. Selleri, *Found. Phys.* **26**, 641–664 (1996).
- [18] J. Bell, in *Speakable and Unspeakable in Quantum Mechanics* (Cambridge University Press, Cambridge, 1988), pp. 67–80.
- [19] S.R. Mainwaring and G.E. Stedman, *Phys. Rev. A* **47**, 3611–3619 (1993).
- [20] C.W. Misner, K.S. Thorne and J.A. Wheeler, *Gravitation* (W.H. Freeman and Company, New-York, 1973), pp. 168–169.
- [21] L.D. Landau and E.M. Lifschitz, *Klassische Feldtheorie* (Akademie-Verlag, Berlin, 1987) p. 297; V.A. Brumberg, *Essential Relativistic Celestial Mechanics* (Adam Hilger, Bristol, 1991), p. 60.
- [22] F. Goy, *Hadr. J.* **20**, Issue 2, 213–218 (1997).
- [23] A. Einstein, *Jahrbuch der Radioaktivität und Elektronik* **4**, 411–462 (1907); **5**, 98–99 (1908) (corrections). The sentence quoted here was translated by myself.
- [24] A.A. Ungar, *Found. Phys.* **26**, 1277–1328 (1996).
- [25] L.H. Thomas, *Nature* **117**, 514 (1926); *Phil. Mag.* **3**, 1–22 (1927).
- [26] J.D. Jackson, *Klassische Elektrodynamik*, 2. verb. Aufl. (De Gruyter, Berlin, 1982).
- [27] V. Bargmann, L. Michel, V.L. Telegdi, *Phys. Rev. Lett.* **2**, 435–436 (1959).
- [28] J. Schwinger, *Am. J. Phys.* **42**, 507–513 (1974).
- [29] R.A. Muller, *Am. J. Phys.* **92**, 313–317 (1992).
- [30] G.P. Fisher, *Am. J. Phys.* **40**, 1772–1781 (1972).
- [31] J. Bailey et al., *Nucl. Phys.* **B150**, 1–75 (1979).
- [32] T. Chang, *Phys. Lett.* **70A**, 1–2 (1979).
- [33] J. Rembieliński, *Phys. Lett.* **78A**, 33–36 (1980).
- [34] F. Selleri, *Found. Phys. Lett.* **9**, 43–60 (1996).